

STIC-ILL

From: STIC-Biotech/ChemLib
Sent: Friday, August 20, 2004 4:18 PM
To: STIC-ILL
Subject: FW: 09705579

508142

NPL	Adonis
MIC	BioTech
NO	Vol NO
Ch Cite	Dupl Request
Call #	

VJ 8/20

-----Original Message-----

From: Yaen, Christopher
Sent: Friday, August 20, 2004 4:05 PM
To: STIC-Biotech/ChemLib
Subject: 09705579

could you please get the following ref(s):

Nouv Rev Fr Hematol. 1989;31(2):77-84.

Cancer Chemother Biol Response Modif. 1991;12:67-73.

Onkologie. 1991 Feb;14(1):7-12.

Drugs. 1992;44 Suppl 4:1-16; discussion 66-9.

Semin Oncol. 1992 Dec;19(6):639-45.

Am J Health Syst Pharm. 1995 Jun 15;52(12):1287-304; quizz 1340-1.

Christopher Yaen
US Patent Office
Art Unit 1642
571-272-0838
REM 3A20
REM 3C18

14828057

Preclinical and Clinical Pharmacology of Vinca Alkaloids

Xiao-Jian Zhou and Roger Rahmani
INSERM U-278, Faculté de Pharmacie, Marseille, France

Summary

Vinca alkaloids, including vinblastine, vincristine, vindesine and vinorelbine, are widely used antineoplastic drugs, either as single agents or in combination with other drugs. The mechanism of action of these cell cycle-dependent agents is the inhibition of tubulin polymerisation into microtubules. Numerous studies have been conducted in animals and humans, using various *in vivo* and *in vitro* models, to investigate the pharmacological behaviour of this class of antitumour drug.

Studies in cellular pharmacology demonstrate that vinca alkaloids are transported by multiple mechanisms, including passive diffusion and energy- and temperature-dependent active transport systems. Moreover, active efflux of drug is involved in the P-glycoprotein-mediated multidrug resistance to vinca alkaloids. This phenomenon may be modulated, *in vivo* and *in vitro*, by calcium antagonists and calmodulin inhibitors.

The clinical pharmacokinetics of vinca alkaloids after intravenous bolus injection, continuous infusion and oral administration are characterised by a large apparent total volume of distribution, high total plasma clearance and long terminal elimination half-life. Biliary excretion is the main elimination pathway, with low urinary excretion. Pharmacokinetic parameters of vinca alkaloids are time- and dose-dependent, and large inter- and intra-individual variabilities have been observed. Human hepatic P-450III_A cytochromes are involved in the metabolism of vindesine, vinblastine and probably other vinca alkaloids. Therefore, the possibility of drug-drug interactions must be considered when coadministering drugs in combination cancer chemotherapy.

Development of newer semisynthetic analogues of vinca alkaloids and conjugation of vinca alkaloids with monoclonal antibodies may result in derivatives with increased antitumour activity and less clinical toxicity.

The early empirical uses of *Catharanthus roseus* (Apocynaceae) for controlling haemorrhage, scurvy, toothache, diabetes, and for the healing of chronic wounds, led to the discovery of antitumour alkaloids in this plant (Johnson et al. 1963). Their anorexic and hypoglycaemic properties prompted isolation and characterisation of the alkaloid molecules responsible for such activities. Further studies then demonstrated that these compounds

produced severe leucopenia by blocking cellular mitosis of haematopoietic tissues.

1. Chemical Structures of the Natural and Semisynthetic Vinca Alkaloids

About 30 alkaloids have been extracted from the periwinkle, of which only vinleurosine, vinrosidine, vinblastine and vincristine possess marked




Fig. 1. Chemical structure of the vinca alkaloid nucleus.

antitumour activity. These compounds, especially vinblastine and vincristine, have been widely used as single agents or in combination with other antineoplastic drugs in cancer chemotherapy for more than 20 years. Besides the naturally occurring alkaloids, some vinca alkaloid analogues have been synthesised by functional transformation (vindesine, desacetylvinblastine-amide) [Cersosimo et al. 1983] or, more recently, by semisynthetic processes (vinorelbine, 5'-noranhydrovinblastine) [Mangne et al. 1979; Maral et al. 1984].

Chemically, these vinca alkaloids have a large dimeric asymmetric structure composed of 2 nuclei linked by a carbon-carbon bond; a dihydroindole nucleus (vindoline), which is the major alkaloid contained in the periwinkle, and an indole nucleus (catharanthine) present in low levels in the plant (fig. 1). The structural differences between vinblastine and vincristine involve the R1 group, while vinblastine and vindesine differ with regard to the R2 and R3 substituents (fig. 2). In contrast with other vinca alkaloids, vinorelbine has structural modifications to the catharanthine nucleus.

2. Mechanism of Action

The mode of action of vinca alkaloids has yet to be completely understood; however, it has been established that antitumour activity is directly related to the high binding affinity of these compounds for tubulin, the basic protein subunit of microtubules. It is generally agreed that these agents arrest cell mitosis at metaphase by preventing tub-

ulin polymerisation to form microtubules and by inducing depolymerisation of microtubules, and are therefore cell cycle-dependent antimitotic agents (spindle poisons). The binding constants of vincristine, vinblastine and vindesine for tubulin are 8.0×10^6 , 6.0×10^6 and 3.3×10^6 mol/L, respectively (Owells et al. 1972, 1977a). The inhibition of net tubulin addition by vinca alkaloids has been evaluated at the assembly ends of bovine brain microtubules. The inhibition constants (K_i) were similar for vincristine, vinblastine and vindesine (0.085, 0.178 and 0.110 μ mol/L, respectively); however, the K_i values correlated poorly with the ability of the substances to inhibit intact cell growth (Jordan et al. 1985). The major difference between these drugs appeared, therefore, to relate to their retention in tumour tissue (Ferguson et al. 1984, 1985; Singer & Himes 1992). It has been suggested that this is conditioned by the formation and stability of vinca alkaloid-tubulin complexes. In mice bearing human rhabdomyosarcoma xenografts with different sensitivities to vincristine, the formation and stability of vincristine-tubulin complexes in cytosols were guanosine 5'-triphosphate (GTP)-dependent. After removal of endogenous GTP, the initial rate of vincristine binding and the maximal level of bound drug were 2- to 3-fold higher in the presence of 0.1 mmol/L GTP than in its absence; the authors suggested that GTP could have had an important role in the therapeutic selectivity of vinca alkaloids (Bowman et al. 1986; Houghton et al. 1987).

2.1 Comparative Effects of Vinca Alkaloids on Axonal Microtubules

Preclinical and clinical studies demonstrated that vinorelbine has a broader antitumour activity with lower toxicity than vinblastine, vincristine and vindesine; this was thought to be because of the action of vinorelbine on microtubules. The work of Binet et al. (1990), using the tectal plate anlage of mouse embryos at the earliest stages of neuronal differentiation, demonstrated that vinblastine, vincristine and vinorelbine induced depolymerisation of interpolar and mitotic microtubules, and cell

Cat




Fig. 2. Structural modifications.

blockage at metaphase concentrations led to pre-centromere microtubule depolymerisation. The active axonal microtubules depolymerised. However, this action served at higher concentrations. Other vinca alkaloids mediated by spiralisation of these data could be a therapeutic index for vincristine neurotoxic effects.

3. Experimental

The cytostatic activity of vinca alkaloids has been studied in established animal and human tumour xenograft models, as well as in tumour xenograft toxicity study using murine cell lines, including murine ovarian carcinoma (A2780) and epidermoid carcinoma (A549).

m microtubules and by of microtubules, and are lent antimitotic agents iding constants of vin- indesine for tubulin are 1.3×10^6 mol/L, re- 1972, 1977a). The inhi- tion by vinca alkaloids ssembly ends of bovine inhibition constants (K_i) e, vinblastine and vin- 0.110 μ mol/L, respec- alues correlated poorly stances to inhibit intact 985). The major differ- appeared, therefore, to umour tissue (Ferguson & Himes 1992). It has conditioned by the for- a alkaloid-tubulin com- human rhabdomyosar- fferent sensitivities to and stability of vincris- cytosols were guanosine ndent. After removal of tial rate of vincristine vel of bound drug were reence of 0.1 mmol/L e authors suggested that portant role in the ther- i alkaloids (Bowman et 1987).

s of Vinca Alkaloids

udies demonstrated that ntitumour activity with astine, vincristine and it to be because of the microtubules. The work g the tectal plate anlage rliest stages of neuronal ed that vinblastine, vin- duced depolymerisation microtubules, and cell

blockage at metaphase. Increasing the drug con- centrations led to progressive depolymerisation of centromere microtubules; however, only vinorel- bine was able to induce complete depolymerisation of these microtubules, resulting in cell arrest at prophase. The activity of these 3 compounds on axonal microtubules was identical; they induced depolymerisation of a labile pool of microtubules. However, this action was dose-related and was ob- served at higher concentrations of vinorelbine than other vinca alkaloids. Since neurotoxicity may be mediated by spirallisation of axonal microtubules, these data could be related to an improved thera- peutic index for vinorelbine, with decreased neu- rotoxic effects.

3. Experimental Antitumour Activity

The cytostatic activity of antitumour vinca al- kaloids has been studied by use of various well es- tablished animal and human tumour cell lines as well as tumour xenografts. A comparative cyto- toxicity study using murine and human tumour cell lines, including murine leukaemia (L1210), human ovarian carcinoma (A2780) and human bronchial epidermoid carcinoma (NSCLC-N6C2), demon-

strated that vinorelbine was as active as vinblas- tine against A2780, less potent than vinblastine and vincristine against L1210, and was more cytotoxic than other vinca alkaloids against NSCLC-N6C2 (Cros et al. 1989; table I).

In a further study using human tumour xeno- grafted *in vivo* in nude mice, vinorelbine was shown to be active against L-27 derived from a human lung carcinoma, whereas vinblastine and vincris- tine were inactive, when all vinca alkaloids were administered intravenously at equitoxic doses [minimal median survival time of treated mice/ median survival time of control mice (T/C) > 0.42]. Similarly, against another human lung carcinoma, LC-06, vinorelbine was more potent than vinblas- tine and vincristine, and was as active as vinde- sine. When human stomach tumours (ST-4 and ST- 40) were tested, vinorelbine displayed greater ac- tivity than other vinca alkaloids (table II).

These *in vitro* and *in vivo* cytotoxicity results have been supported by clinical application of vinca alkaloids. Indeed, vincristine and vindesine are generally used in the treatment of various leukaemias, whereas the main clinical indication for vi- norelbine is the treatment of non-small cell lung cancer.




Fig. 2. Structural modifications to the vindoline and catharanthine nuclei in various vinca alkaloids.

Table I. Comparison of cytotoxicity of vinca alkaloids against murine and human tumour cell lines (after Cros et al. 1989)

Cell lines	IC ₅₀ (nmol/L)			
	vinorelbine	vinblastine	vincristine	vindesine
Human ovarian carcinoma (A2780)	47.0	42.6	ND	ND
Murine leukaemia (L1210)	7.2	5.0	3.4	ND
Human bronchial epidermoid carcinoma (NSCLC-N6L2)	1.4	2.6	36.0	2.8

Abbreviations: IC₅₀ = concentration inhibiting 50% of cell growth; ND = not determined; NSCLC = non-small cell lung cancer.

Table II. Comparative effect of vinca alkaloids on human tumour xenografts (after Cros et al. 1989)

Tumour	line	Minimal T/C (%)			
		vinorelbine	vinblastine	vincristine	vindesine
Lung	L-27	0.27	0.48	ND	0.52
Lung	LC-06	0.06	0.25	0.27	0.06
Stomach	ST-4	0.28	0.35	0.44	0.32
Stomach	ST-40	0.21	ND	ND	0.39

Abbreviations: T/C = median survival time of treated mice (T) divided by median survival time of control mice (C); ND = not determined.

4. Mechanism of Transport

Numerous *in vitro* studies have been conducted with animal or neoplastic cells to elucidate the mechanism of transport of vinca alkaloids. Irrespective of the model used, they appeared to be intensely and rapidly taken up into cells; however, there were marked differences between the alkaloids. In suspensions of freshly isolated rat hepatocytes, vinblastine was more intensively accumulated in the cells than vincristine or vindesine (Rahmani et al. 1990; Zhou et al. 1990). Similar results were obtained using human hepatocytes, in which the intensity of cellular accumulation of vinca alkaloids appeared to be directly proportional to their liposolubility. Moreover, the intracellular drug concentration was much higher than the extracellular one (Cano et al. 1988). In murine leukaemia cells, the intracellular vincristine concentration was 5- to 20-fold higher than the extracellular concentration (Bleyer et al. 1975). Similar results have been reported with murine lymphoma cell lines (Heja and S49) and a human leukaemic

cell line (HL 60), the ratio of intracellular to extracellular drug concentrations varying from 150 to 500 for vinca alkaloids (Ferguson et al. 1984, 1985).

Multiple mechanisms have been described for the transmembrane passage of vinca alkaloids. Some authors have reported an energy- and temperature-dependent active transport; the Michaelis-Menten transport constant (K_m) for vincristine is 6.45 $\mu\text{mol/L}$ in a human leukaemic cell line (CEM/CCRI) and 9.2 $\mu\text{mol/L}$ in murine leukaemic cell lines (L1210, P388) [Beck 1983; Bleyer et al. 1975]. Our own results, obtained by using purified plasma membrane vesicles prepared from rat liver, demonstrated that vinorelbine-binding abilities (from 0.01 to 0.03 $\mu\text{mol/L}$) accounted for 39 to 48% of total uptake. Transport of vinorelbine was intense and rapid, and primarily occurred by a temperature-independent and unsaturable process (consistent with a simple diffusion mechanism), which accounted for the majority of transportation. A secondary mechanism involved a temperature-dependent saturable transport system. This process was in accord with Michaelis-Menten ki-

netics, exhibiting 3 binding sites ($K_{d1} = 0.1 \mu\text{mol/L}$, $K_{d3} = 55 \mu\text{mol/L}$) [Rahmani-Jourdeuil et al. 1990].

5. In Vitro Biotransformation

The metabolism of vinca alkaloids was studied *in vitro* using animal hepatocytes in subcellular fractions such as microsomes, incubated in suspensions of microsomes. Vincristine, vinorelbine and vindesine were completely converted to metabolites which were rapidly excreted from the medium. The intracellular accumulation of most exclusively unchanged drug of total intracellular drug was due to tubulin protein (Rahmani 1990). A large variability of results was observed when incubated with hepatic microsomes of different species (rat, rabbit, human). Similarly, by using human hepatic microsomes, the biotransformation of vinca alkaloids presented individual variability (fig. 4) which explained the clinical pharmacokinetics of these drugs (Zhou et al. 1990). This bank of microsomes




Fig. 3. Scatchard analysis of vinorelbine binding to microsomes showing that there are 3 different binding sites (1, 2, 3) associated with 3 different microsomes (Rahmani-Jourdeuil et al. 1990).

ter Cros et al. 1989)

a videsine

ND

ND

2.8

on-small cell lung cancer.

e videsine

0.52

0.06

0.32

0.39

vide (C); ND = not determined.

netics, exhibiting 3 heterogeneous high-affinity binding sites ($K_{d1} = 0.1 \mu\text{mol/L}$, $K_{d2} = 16.5 \mu\text{mol/L}$, $K_{d3} = 55 \mu\text{mol/L}$) [fig. 3].

5. In Vitro Biotransformation

The metabolism of vinca alkaloids has been studied *in vitro* using freshly isolated human and animal hepatocytes in suspension, and hepatic subcellular fractions such as microsomes. When incubated in suspensions of rat hepatocytes, vinblastine, vincristine and vindesine were almost completely converted into several metabolites, which were rapidly excreted into the extracellular medium. The intracellular medium contained almost exclusively unchanged drug (more than 85% of total intracellular drug), presumably bound to tubulin protein (Rahmani et al. 1990; Zhou et al. 1990). A large variability in vinca alkaloid metabolism was observed when these compounds were incubated with hepatic microsomal fractions from different species (rat, rabbit, dog, pig, baboon, and human). Similarly, by using a bank of 30 different human hepatic microsomes, the biotransformation of vinca alkaloids presented significant interindividual variability (fig. 4), which may partially explain the clinical pharmacokinetic characteristics of these drugs (Zhou et al. 1992). Recently, using this bank of microsomes, we demonstrated that

human hepatic P-450III_A cytochromes were predominantly involved in the biotransformation of vindesine and vinblastine, and probably all vinca alkaloids.

Since vinca alkaloids are frequently administered in combination with other antineoplastic drugs, potential metabolic drug interactions have been evaluated. Results indicated that some drug classes such as the epipodophyllotoxins (e.g. etoposide, teniposide) significantly inhibited vinca alkaloid metabolism (Zhou et al. 1992). These findings will aid optimisation of combination chemotherapy protocols.

6. Multidrug Resistance

The clinical use of vinca alkaloids has led to the development of resistant tumour cells, which has limited effective cancer chemotherapy with these agents. Furthermore, cells selected for resistance to vinca alkaloids also exhibit cross-resistance to a variety of other structurally unrelated 'natural' compounds such as anthracyclines and epipodophyllotoxins, despite no previous exposure to these drugs (Beck-Hansen et al. 1976; Carlsen et al. 1976). This is a major feature of the phenomenon known as pleiotropic or multidrug resistance (MDR) (Beck 1983; Ling et al. 1983), which appears to result from decreased intracellular drug retention, due to an enhanced efflux (Dalton et al. 1986; Dano 1973; Inaba & Johnson 1977). For a given extracellular drug concentration, multidrug-resistant cells maintain a lower intracellular drug concentration than sensitive parental cells.

Numerous studies have been done to determine the origin of MDR by detecting biochemical alterations in multidrug-resistant cells compared with intact cells. The most consistent change is the overexpression of a high molecular weight (170kD) surface plasma membrane glycoprotein, named P-glycoprotein (Gp-170) in multidrug-resistant cells (Beck et al. 1979; Biedler & Peterson 1981; Bosmann 1971; Juliano & Ling 1976; Kartner et al. 1983). In vinblastine-resistant human leukaemia cells, a surface glycoprotein with a molecular weight of 180 to 210kD was detected by use of mono-




Fig. 3. Scatchard analysis of the vinorelbine transport systems showing that there are 3 binding sites (marked 1, 2 and 3) associated with 3 different dissociation constants (after Rahmani-Jourdeuil et al. 1991).




Fig. 4. Interindividual variability in vindesine metabolism on human hepatic microsomal fractions (after Zhou et al. 1992).

clonal antibodies (Danks et al. 1985). The P-glycoprotein was found to be a receptor for vinca alkaloids, and tumour cell drug resistance appeared to be proportional to the amount of P-glycoprotein. Structural analysis of P-glycoprotein demonstrated that it is a conserved integral plasma membrane protein with structural features characteristic of an energy-dependent plasma membrane transport system (Fry et al. 1986; Walker et al. 1982). Moreover, it has been suggested that P-glycoprotein binds adenosine triphosphate (ATP) and couples ATP hydrolysis to its respective energy-requiring biological processes (Higgins et al. 1986). Therefore, one of the dominant mechanisms for resistance to vinca alkaloids or other drug classes could be the reduced intracellular drug retention after P-glycoprotein-mediated, energy-dependent, drug efflux. However, some multidrug-resistant phenotypes have appeared to be P-glycoprotein-independent. Vinca alkaloid-resistant human leukaemic lymphoblast cells (CCRI-CEM) treated with pronase or tunicamycin (an inhibitor of P-glyco-

protein), which resulted in the absence of resistance-associated glycoprotein, continued to express resistance to these drugs (Beck & Cirtain 1982). Moreover, therapy-induced resistance of a human leukaemic cell line (LALW-2) to vinca alkaloids has also been shown to be independent of P-glycoprotein (Haber et al. 1989). Gene amplification appears to be the basis for increased expression of P-glycoprotein in highly resistant cell lines. The amplified DNA detected in multidrug-resistant cell lines contains the P-glycoprotein multigene family (*mdr1* gene) [Van der Bliek et al. 1986]. Usually, only 1 subset of the *mdr1* gene members is highly amplified in different multidrug-resistant cell lines (De Bruijn et al. 1986; Riordan et al. 1985). Two of this gene family, *mdr1a* and *mdr1b*, encode the 120 and 125kD P-glycoprotein precursors, respectively, in the multidrug-resistant murine J744.2 cell line. The P-glycoprotein encoded by *mdr1a* is functionally more efficient than the form encoded by *mdr1b*, either of which may be overexpressed in multidrug-resistant cells (Lothstein et al. 1989).

Clinically, MDR is prevented by combination drugs to which the resistant. Since tumours exhibit low cross-resistance and other vinca alkaloids limit the development of resistant lines (Maral et al. 1991), published data on MDR focused on screening metabolic activity or toxicity by cellular or tumoral model development of the mechanism is directly related to the surface P-glycoprotein, with cellular drugs, many can overcome resistance at agents such as calcium inhibitors. Verapamil, coronary vasodilator agent, is potent at enhancing vincristine *in vitro* and verapamil to overcome MDR in multidrug-resistant murine cells. Vincristine and vinblastine were greatly increased by verapamil (Tsuruo et al. 1988) to be greater for vinca alkaloids than 'natural' drugs such as dophyllotoxins (Beck et al. 1982).

Calmodulin is a regulatory protein involved in the assembly-disassembly process of the microtubule, which is the most important target of the vinca alkaloids (Beck et al. 1979). Therefore it is suggested that the vinca alkaloid-resistant tumour cells may be more sensitive to microtubule inhibitors, prenylamine and ipramine. In the presence of 10 μ mol/L, vincristine cytotoxicity was 10-fold in vincristine-resistant (K562) cells. The accumulation appeared to be an enhancement of vincristine cytotoxicity; however, it did not correlate with the cytotoxic effect.



ons (after Zhou et al. 1992).

n the absence of resist-
in, continued to express
(Beck & Cirtain 1982).
d resistance of a human
-2) to vinca alkaloids has
dependent of P-glycopro-
Gene amplification ap-
increased expression of
sistant cell lines. The am-
multidrug-resistant cell
protein multigene family
ck et al. 1986]. Usually,
gene members is highly
tidrug-resistant cell lines
Jordan et al. 1985). Two
a and *mdr1b*, encode the
rotein precursors, respec-
sistant murine J744.2 cell
ncoded by *mdr1a* is func-
an the form encoded by
may be overexpressed in
Lothstein et al. 1989).

Clinically, MDR has generally been circumvented by combination therapy or by the use of drugs to which the cancer cells are not cross-resistant. Since tumour cell lines have been shown to exhibit low cross-resistance between vinorelbine and other vinca alkaloids, the use of this drug may limit the development of multidrug-resistant cell lines (Marai et al. 1981). So far, however, published data on MDR reversal have essentially focused on screening molecules without antitumour activity or toxicity by means of experimental cellular or tumoral models. Since, in most cases, the development of the multidrug-resistant phenotype is directly related to the overexpression of the surface P-glycoprotein, which actively extrudes intracellular drugs, many of the molecules that could overcome resistance are cellular membrane active agents such as calcium antagonists and calmodulin inhibitors. Verapamil, a calcium antagonist with coronary vasodilator activity, is known to be very potent at enhancing vinca alkaloid cellular accumulation *in vitro* and *in vivo*. The ability of verapamil to overcome MDR was first observed in multidrug-resistant murine P388 cell lines, in which vincristine and vinblastine intracellular concentrations were greatly increased in the presence of verapamil (Tsuruo et al. 1981). The effect was found to be greater for vinca alkaloids than for the other 'natural' drugs such as anthracyclines and epipodophyllotoxins (Beck et al. 1986).

Calmodulin is a regulatory protein of the assembly-disassembly processes of microtubules, which are formed by the polymerisation of tubulin, the most important target of vinca alkaloids (Welsh et al. 1979). Therefore it is reasonable to expect that the vinca alkaloid-resistance of cell lines or human tumour cells may be modulated by the calmodulin inhibitors, prenylamine, trifluoperazine, and clomipramine. In the presence of these agents at 6.6 μ mol/L, vincristine cytotoxicity increased 29- to 45-fold in vincristine-resistant human erythroleukemia (K562) cells. The enhancement of vincristine accumulation appeared to be related to the enhancement of vincristine cytotoxicity by these agents; however, it did not always correlate directly with the cytotoxic effect. Trifluoperazine and

clomipramine led to high vincristine accumulation in vincristine-resistant K562 cells, while prenylamine induced a lower accumulation of vincristine. However, prenylamine caused greater enhancement of vincristine cytotoxicity (Tsuruo et al. 1983). A number of other drugs, including flunarizine, reserpine, vitamin A, cepharantine, cefoperazone, quinidine, cyclosporin and its analogues, hexamethylene bisacetamide and some vinca alkaloids without antitumour activity, have also been reported to reverse vinca alkaloid-related multidrug-resistant phenotypes (Ford & Hait 1990).

The mechanism of these drugs in overcoming MDR is not fully elucidated. However, verapamil, trifluoperazine, quinidine, reserpine and cyclosporin share some or all of the features of the known substrates of the P-glycoprotein, including lipophilicity, a planar polycyclic stereochemistry and weak basic properties. These agents appear to reduce or overcome the multidrug resistant phenotype by acting as substrates for the active efflux pump mediated by the P-glycoprotein, competitively inhibiting the active efflux of cytotoxic drugs, thereby increasing intracellular drug accumulation and cytotoxicity. However, it appears necessary to maintain constant intracellular concentrations for the entire time during which cells are exposed to the cytotoxic agent. The calcium antagonists have proven more active than the calmodulin inhibitors in circumvention of MDR. To date, these agents have not been used successfully in clinical practice, since the concentrations required to overcome MDR *in vitro* often lead to severe side effects (such as cardiac disturbances) *in vivo*.

7. Pharmacokinetic Properties of Vinca Alkaloids

7.1 Analytical Methods

Pharmacokinetic studies of vinca alkaloids were initially conducted with radiolabelled drugs, and their concentrations in biological fluids calculated according to the radioactivity; however, this technique could not distinguish unchanged drug from metabolites and/or degradation products, making the results less reliable than currently available as-

say techniques. Radioimmunoassay (RIA) and enzyme-linked immunosorbent assay (ELISA) methods have greatly facilitated the clinical pharmacokinetic investigation of these compounds and have enabled their re-evaluation. These immunological assay methods present various advantages with regard to specificity, sensitivity, and reproducibility, so that they are now the standard assay methods used.

RIA methods using [125 I]-labelled probes as antigen have been successfully applied in vindesine and vinorelbine pharmacokinetic evaluations (Rahmani et al. 1983, 1984a). A highly specific RIA for vincristine without cross-reactivity for vinblastine exists (Huhtikangas et al. 1987), and a new ELISA method with a detection limit of 5 nmol/L has been used to re-evaluate the clinical pharmacokinetics of vinblastine (Hacker et al. 1984). In addition, antibodies used for RIA were generally produced in rabbits, but now more specific and sensitive antibodies may be produced using monoclonal techniques (Pontarotti et al. 1985).

Other techniques have been developed, including high performance liquid chromatography (HPLC) [Bloemhof et al. 1991; De Smet et al. 1985; Jehl et al. 1990; Vendrig et al. 1988a,b], which is able to separate unchanged drug from its metabolites. Advances in the field of extraction (solid phase) and detection (electrochemical and fluorescence) techniques make this method applicable for clinical pharmacokinetic studies of vinca alkaloids.

7.2 Clinical Pharmacokinetics

In cancer chemotherapy, the vinca alkaloids have usually been administered by direct intravenous injection at doses of 1 to 2, 1.5 to 6, 7 to 11, and 15 to 30 mg/m² for vincristine, vinblastine, vindesine, and vinorelbine, respectively. The pharmacokinetics of vinca alkaloids can generally be described by an open 3-compartment model, which was used to derive pharmacokinetic parameters (Rahmani et al. 1988; fig. 5). The clinical pharmacokinetics of vinca alkaloids have been characterised by a large volume of distribution, a high systemic clearance, and a long terminal eliminat-





Fig. 5. Plasma pharmacokinetics of vindesine (8 mg/m²; ○) and vinorelbine (30 mg/m²; ●) in patients after intravenous injection (after Rahmani et al. 1988).

ion half-life with large inter-molecule and inter- and intra-individual variability in pharmacokinetic parameters. While the initial and intermediate phase half-lives of these agents were similar after intravenous injection, there was a marked difference in the apparent terminal-phase half-lives: 85 hours for vincristine, about 24 hours for vinblastine and vindesine (Nelson et al. 1982) and 40 hours for vinorelbine (Rahmani et al. 1986, 1987) [table III]. The long terminal elimination half-life of vincristine associated with its lower elimination constant may explain its lower maximum tolerated dose compared with other vinca alkaloids. Moreover, the percentages of central compartment distribution volume were higher for vincristine (33%) and vinblastine (69%) than for vindesine (5.4%), suggesting that the latter binds less rapidly to formed blood elements than vincristine and vinblastine (Nelson et al. 1982). Similarly, the relatively high distribution volume of these compounds reflects their large tissue distribution (table III).

The vinca alkaloids are cell cycle-dependent antimitotic agents with narrow therapeutic windows. Intravenous injection leads to very high initial plasma drug concentrations (several hundred µg/L), which may cause more toxic than therapeutic effects. In comparison with intravenous injection,

Table III. Pharmacokinetic parameters of vinca alkaloids in patients after intravenous injection

Dose (mg/m ²)	Vinblastine			Vincristine			Vindesine			Vinorelbine		
	RIA	RIA	RIA	RIA	RIA	RIA	RIA	RIA	RIA	RIA	RIA	HPLC
$t_{1/2\alpha}$ (h)	0.065	7-14	3	0.056	2	1.5-4	2.5	0.4-8	15-30	30	0.067	0.134
$t_{1/2\beta}$ (h)	0.86	1.64	2.58	2.27	85.0	20.2	0.912	22.7	1.82	79.8	1.82	2.08
$t_{1/2\gamma}$ (h)	19.5	24.8	85.0	85.0	85.0	20.2	24.2	22.7	35.0	31.2	35.0	42.0



Plasma concentration of vindesine (8 mg/m²; O) in patients after intravenous injection (1988).

inter-molecule and inter-patient variability in pharmacokinetic parameters was similar after intravenous injection. There was a marked difference in terminal-phase half-lives: 85 hours at 24 hours for vinblastine (Rahmani et al. 1982) and 40 hours for vindesine (Rahmani et al. 1986, 1987) [table 1]. The elimination half-life of vindesine is lower than that of vinblastine. The maximum tolerated dose of vinca alkaloids. More variability in central compartment distribution for vincristine (33%) than for vindesine (5.4%). Vindesine binds less rapidly to tubulin than vincristine and vinblastine (Rahmani et al. 1982). Similarly, the terminal volume of distribution of vindesine is larger than that of vinblastine.

Vindesine is cell cycle-dependent and has a narrow therapeutic window. The initial plasma concentrations (several hundred µg/L) are toxic than therapeutic after intravenous injection.

Table III. Pharmacokinetic parameters of vinca alkaloids in patients after intravenous injection

	Vinblastine			Vincristine			Vindesine			Vinorelbine			HPLC
	RIA	RIA	RIA	RIA	RIA	RIA	RIA	RIA	RIA	RIA	RIA	RIA	
Dose (mg/m²)	0.065	0.062	7-14	3	0.056	0.077	1.5-4	0.054	0.037	0.4-8	15-30	30	30
t _{1/2} α1 (h)	0.88	1.64		2.58	0.056	0.077	0.054	0.037	0.037	0.067	0.067	0.134	0.134
t _{1/2} α2 (h)					2.58	2.27	1.65	0.912	0.912	1.82	1.82	2.08	2.08
t _{1/2} γ (h)						85.0	20.2	24.2	22.7	31.2	35.0	79.8	42.0
V _c (L/kg)	0.28	0.69			0.33	0.33	0.084	0.054	0.054				
V _z (L/kg)	25.2	27.3			8.4	8.4	11.2	8.8			26.9	75.6	75.6
CL (L/h/kg)						0.106	13.2			0.519	0.920	0.418	1.26
Ae (% of total dose)				9.5					4.2		3.8	24.6	10.9
Reference no.	1	2		1	2	2	1	2	2	3	4	5	6

Abbreviations: RIA = radioimmunoassay; HPLC = high performance liquid chromatography; t_{1/2}α1 = half-life of first phase; t_{1/2}α2 = half-life of second phase; t_{1/2}γ = terminal elimination half-life; V_c = apparent volume of distribution of central compartment; V_z = apparent volume of distribution during the terminal phase; CL = clearance of drug from plasma; Ae = amount of unchanged drug excreted into urine.
Reference key: 1 Owles et al. (1977b); 2 Nelson et al. (1982); 3 Rahmani et al. (1984b); 4 Rahmani et al. (1986); 5 Rahmani et al. (1987); 6 Boré et al. (1989); 7 Jerni et al. 1991.

continuous infusion may be advantageous, because it avoids the toxic peak concentrations and may increase the duration of exposure of cells to the effective concentration (Brade 1981). The steady-state plasma concentration after infusion of vinca alkaloids is in the range of a few µg/L, which seems to be effective because of the high affinity of these agents for tubulin. Increased antitumour activity, with little or no increase in toxicity after continuous infusion of vinca alkaloids has been reported: high remission rates resulted from long-term infusion of vindesine or vinorelbine in patients with haematological malignancies, breast cancer, or head and neck cancer, even when bolus injection had failed (table IV; Bodey et al. 1980; Hande et al. 1980; Mathé et al. 1981). Steady-state levels of 6 to 15 µg/L were achieved with vindesine (Rahmani et al. 1985); this concentration has been shown to produce cytotoxic effects on mammalian cell cultures. For vinblastine, steady-state plasma concentrations of 1.5 to 2 µg/L prevented severe myelosuppression (Ratain & Vogelzang 1986). The dosages administered by continuous infusion were 0.5 to 1, 1.2 to 2, and 0.8 to 1.5 mg/m²/day for vincristine, vinblastine, and vindesine, respectively. In addition, a significant circadian variation in the plasma vindesine concentration was observed during a 48-hour 1.5 mg/m²/day continuous infusion in 9 patients (Focan et al. 1989). This finding is of importance with regard to schedule optimisation for drugs with a long terminal half-life. The systemic clearance of vincristine (Jackson et al. 1981), vinblastine (Lu et al. 1983; Ratain et al. 1987) and vindesine (Rahmani et al. 1985) has been reported to decrease during the infusion period compared with intravenous injection. This schedule-dependent systemic clearance could be explained by nonlinear pharmacokinetics, which appear to be a general feature of vinca alkaloids; nonlinearity was strongly suggested by the apparent dose- and/or time-dependence of vindesine and vinorelbine pharmacokinetics after bolus doses (Rahmani et al. 1984b, 1986).

All available data show a large intra- and inter-patient variability in pharmacokinetic parameters for vinca alkaloids. This could result from indi-

Table IV. Comparison of response rates and toxicities of bolus injection versus continuous infusion of vindesine

	Refractory breast cancer ¹			Pretreated melanoma, NSCLC ²			Leukaemia, haematosarcoma ³		
	bolus	infusion		bolus	infusion		bolus	infusion	infusion
Doses (mg/m ² /day)	3-5 every 10d	1.0-1.2 x 5d 1.4-1.5 x 5d every 3 weeks		3.0 x 2d every 2 weeks	3.0 every week		2.0 x 2d every week	2.0 x 2d after failure of bolus	2.0 x 2d as first treatment
No. of responses/no. of patients treated									
Complete response	1/23	6/21		1/17			10/43	2/12	0/3
Partial remission	4/23	10/21		2/17	1/9		6/43	4/12	1/3
No cure									
No. of patients with toxicity/no. of patients treated									
Paraesthesia	13/52	14/110		8/17	0.9				
Reflexes decreased	13/52	7/110							
Muscle weakness	4/52	7/110							
Haematology									
WBC ($\times 10^9/L$) ^a	2.6-3.3	1.5-2.0		< 3.0 (8/17) ^b	< 3.0 (4/9) ^b				
Platelets ($\times 10^9/L$) ^a	173-200	198-201		< 100 (0/9) ^b	< 100 (1/17) ^b				

^a Values are given as ranges.^b No. of patients with toxicity/no. of patients treated.

Abbreviations: NSCLC = non-small cell lung cancer; WBC = white blood cell.

Reference key: 1 Bodey et al. (1980); 2 Hande et al. (1980); 3 Mathé et al. (1981).

vidual differences in metabolism (Jackson et al. 1982) and uncertainties observed in the dependencies observed (Rahmani et al. 1986, 1987); of 24 patients treated with bolus injection followed by precontinuous infusion, the serum albumin levels and bolus clearance ($r = 0$) and clearance ($r = 0.39$). If bolus clearance was albumin and negatively correlated (Nelson et al. 1987). Moreover, vincristine, vinblastine and vinorelbine be significantly correlated (Nelson et al. 1982) as a cause of this, some other relationships between pharmacokinetic parameters and hydro modification or ring increases clearance of substituent on the vinyl and increases the toxicity (Rahmani et al. 1986). The tumour activities, are also correlated (Maral et al. 1986) [table V].

The large volume of distribution and its high affinity for continuous exposure to low

Table V. Relationship between

Vinca alkaloids

Vinblastine

Vincristine

Vinorelbine

Sodium-formyl vinorelbine

Desacetyl-vinorelbineamide

Abbreviations: LD₁₀ = dose

Reference key: 1 Rahmani et al.

vidual differences in hepatic drug disposition and metabolism (Jackson et al. 1981; Van den Berg et al. 1982) and undefined time- and/or dose-dependencies observed after administration (Rahmani et al. 1986, 1987; Zhou et al. 1991). In a group of 24 patients treated with vinblastine by bolus injection followed by prolonged (for 2 to 36 weeks) continuous infusion, the interpatient differences in serum albumin levels were correlated with both the bolus clearance ($r = 0.49$) and the initial infusion clearance ($r = 0.39$). Inpatient variation in vinblastine clearance was positively correlated with albumin and negatively correlated with dose (Ratain et al. 1987). Moreover, the systemic clearance of vincristine, vinblastine and vindesine was found to be significantly correlated with weekly dosages (Nelson et al. 1982) and to parallel toxicity. Because of this, some authors have tried to establish relationships between chemical structure and pharmacokinetic parameters. In general, the 5'-noranhydro modification of the original catharanthine ring increases clearance, while changes to a formyl substituent on the vindoline ring reduces clearance and increases the terminal elimination half-life (Rahmani et al. 1986). Toxicities, and possibly antitumour activities, are related to changes in clearance (Maral et al. 1981, 1984; Todd et al. 1976) [table V].

The large volume of distribution of vinorelbine and its high affinity for tubulin suggest that continuous exposure to low concentrations may be ef-

fective and could easily be achieved by oral administration. Recent results obtained in phase I studies of this drug demonstrated that oral vinorelbine was rapidly absorbed and had a good bioavailability (about 40%) [Rahmani et al. 1991]. The relatively high bioavailability could be related to its high liposolubility. Moreover, oral administration of vinorelbine resulted in a pharmacokinetic profile and antitumour activity that were similar to those with intravenous injection, and could therefore represent a new drug delivery pathway for antitumour vinca alkaloids in ambulatory cancer chemotherapy (Zhou et al. 1991).

In human subjects, the vinca alkaloids are largely metabolised and eliminated via the hepatobiliary system. In 2 patients treated with [^3H]-vinorelbine, Boré et al. (1989) reported the presence of circulating metabolites, as shown by large discrepancies between plasma concentrations determined by RIA and direct counts of radioactivity: results from one patient are shown in figure 6. Faecal drug excretion within 21 days represented 46.2% of total dose, while urinary excretion was 24.6%. Urinary excretion of vinorelbine was, however, slightly higher than with other vinca alkaloids; urinary excretion was reported to be about 10% of the total dose for vinblastine and vindesine (Rahmani et al. 1984b). However, only desacetyl vinblastine (Owells et al. 1977b) and desacetyl vinorelbine (Jehl et al. 1991) have been identified as urinary metabolites of vinblastine and vinorelbine, respectively.

Table V. Relationship between clearance and toxicity of vinca alkaloids in animal models

Vinca alkaloids	Mean (\pm SD) clearance (L/kg \cdot h) ¹	Mean (\pm SD; LD ₅₀ (mg/kg)		Mean LD ₁₀ (mg/kg) ³
		rat	mouse	
Vinblastine	1.2 \pm 0.4	2.9 \pm 1.5 ²	10.3 \pm 0.8 ²	10.0
Vincristine	0.13 \pm 0.5	1.0 \pm 0.1 ²	2.1 \pm 0.1 ²	1.4
Vinorelbine	2.1 \pm 0.5		24.0 ³	20.0
Sodium-formyl vinorelbine	0.27 \pm 0.14		5.0 ³	
Desacetyl-vinorelbinamide	2.4 \pm 0.7		17.0 ³	

Abbreviations: LD₁₀ = dose resulting in death of 10% of animals; LD₅₀ = median lethal dose.

Reference key: 1 Rahmani et al. 1986; 2 Todd et al. 1976; 3 Maral et al. 1984.

a. Values are given as ranges.
b. No. of patients with toxicity/no. of patients treated.
Abbreviations: NSCLC = non-small cell lung cancer; WBC = white blood cell.
Reference key: 1 Bodey et al. (1980); 2 Hande et al. (1980); 3 Mathis et al. (1981).




Fig. 6. Plasma pharmacokinetics of vinorelbine in a patient after the administration of 30 mg/m² (49.5mg) of [³H]-vinorelbine: evidence for the presence of circulating vinorelbine-related metabolites; ● = direct counts of radioactivity; ▲ = radioimmunoassay (after Boré et al. 1989).

8. Clinical Efficacy and Tolerability of Vinca Alkaloids

8.1 Efficacy

Significant differences have been observed in the antitumour activity and toxicity of vinca alkaloids. Vinblastine is usually used to treat Hodgkin's and non-Hodgkin's lymphomas and some solid tumours, including testicular and breast cancer. Vincristine exhibits substantial activity against Hodgkin's and non-Hodgkin's lymphomas, acute lymphoblastic leukaemia, breast carcinoma, Wilm's tumour, Ewing's sarcoma, neuroblastoma, hepatoblastoma and embryonal rhabdomyosarcoma. The antitumour activity of vindesine was found to be similar to that of vinblastine and vincristine, and it is generally used in the treatment of childhood acute lymphocytic leukaemia, acute granulocytic leukaemia, small cell lung cancer and non-Hodgkin's lymphomas. Clinical trials demonstrated that vinorelbine was highly effective in at least 4 categories of cancer: non-small cell lung cancer, breast cancer, ovarian cancer and Hodgkin's disease.

In addition, it has been reported that vinca alkaloid administration led to high platelet counts (Retsas et al. 1978), which appeared to be partic-

ular to the alkaloids and was used with success in the management of idiopathic thrombocytopenic purpura and autoimmune thrombocytopenia.

8.2 Tolerability

The toxicity of vinca alkaloids has been extensively studied. Vincristine administration has resulted in disturbances of the central, peripheral and autonomic nervous systems, characterised by a decrease in deep tendon reflexes, paraesthesias, constipation, myalgias, muscle weakness and paralytic ileus (Brade 1981; Legha 1986). Administration of vinblastine and vindesine has led mainly to haematological toxicities, including anaemia and leucopenia (Cersosimo et al. 1983). In addition, these agents have commonly induced alopecia and gastrointestinal disturbances. With vinorelbine, neurotoxicity has been very mild and generally limited to a decrease in or abolition of the osteotendinous reflex. Paraesthesia and cases of paralytic ileus have been rare with this agent. The major dose-limiting factors are leucopenia for vinblastine, vindesine and vinorelbine, and neurotoxicity for vincristine. The neurotoxic symptoms induced by these agents appear to be dose- and time-dependent and generally resolve with dose reduction or treatment withdrawal.

9. Analogue Development

Efforts continue to be devoted to the development of new analogues of vinca alkaloids. Vinzolidine is a semisynthetic analogue of vinblastine. Given orally, this compound has marked cytotoxicity against lymphoma cells; however, phase II trials revealed side effects such as severe bone marrow toxicity (Budman et al. 1984). The coupling of vinca alkaloids with amino acids has led to some interesting analogues. Vinca-23-oyl amino acid derivatives, *L*-tryptophan-*o*-esters and deoxy-*o*-esters were found to have superior chemotherapeutic activity and diminished toxicity compared with the parent compound in experimental systems (Rao et al. 1985). More recently, 2 vinblastine analogues, S 12363 and S 12362, have been synthesised by

grafting an optically active amino acid at the C23 position of 12363 was as potent as vinblastine in *in vitro* tests; however, 2 murine and 6 human tumours were found to be 7- to 553-fold more cytotoxic than vincristine. S 12362, which has the configuration of the side chain atom of the side chain compared with its *in vivo* activity. Intravenously or intraperitoneally, S 12362 was at least as active as vinblastine while the optimal dose was 0.15 to 0.20 mg/kg vs 0.15 to 0.20 mg/kg for vinblastine. Its high potency was due to its properties compared with vinblastine, such as its ability to enter cellular retention.

Monoclonal antibody conjugates have gained as potentially useful tools for the treatment of human cancers. It is possible to specifically bind to surface antigens, which are attached to cytotoxic agents. A number of monoclonal antibodies have been synthesised both *in vitro* and *in vivo*. One example is the adenocarcinoma cell line 4S2, coupled through its 4-hydroxy group of the succinate bridge, compared with vinblastine molecules. It inhibited a significant amount of human lung carcinoma xenografts in mice (1987). Pharmacokinetic studies in monkeys indicated that the conjugate was mostly in the intact form, not catabolised in the liver or excreted in the urine of vinca alkaloids (et al. 1987). Recently,

as used with success in
thrombocytopenic
thrombocytopenia.

caloids has been exten-
administration has re-
central, peripheral and
characterised by a de-
xes, paraesthesias, con-
weakness and paralytic
(1986). Administration of
has led mainly to hae-
molytic anaemia and leu-
kopenia (1983). In addition, these
agents caused alopecia and gas-
trointestinal toxicity. With vinorelbine, neu-
ropathy and generally limited
toxicity of the osteotendinous
tissues. Side effects of paralytic ileus have
been the major dose-limiting
toxicities of vinorelbine, vindesine and
vinorelbine. The
toxicity caused by these agents
is dose-dependent and gener-
ally reversible after treatment

ient

devoted to the develop-
ment of vinca alkaloids. Vinorelbine
is a derivative of vinblastine.
It has marked cyto-
toxicity; however, phase II
studies as severe bone mar-
row suppression (1984). The coupling of
vinorelbine to antibodies has led to some
improvement in the activity of
vinorelbine. 23-oxo amino acid de-
rivatives and deoxy-o-esters
of vinorelbine have been synthesised by

grafting an optically active α -aminophosphonate at
the C23 position of 04-desacetyl vinblastine. S
12363 was as potent as vincristine, and less potent
than vinblastine in inhibiting tubulin polymerisa-
tion *in vitro*; however, when tested on a panel of
2 murine and 6 human tumour cell lines, it was
found to be 7- to 553-fold and 12- to 74-fold more
cytotoxic than vincristine and vinblastine, respec-
tively. S 12362, which differs from S 12363 only
by the configuration of the asymmetrical carbon
atom of the side chain, was 18- to 59-fold less toxic
compared with its isomer. When administered
intravenously or intraperitoneally in mice bearing
P388 ascites tumour and B16 melanoma, S 12363
was at least as active as vincristine and vinblastine,
while the optimal dose was 10- to 20-fold lower
(0.15 to 0.20 mg/kg vs 2 to 5 mg/kg). Thus, S 12363
has been considered to be the most potent deriva-
tive. Its high potency could in part be accounted
for by properties conferred by the α -aminophos-
phonic acid, such as facilitated uptake and/or bet-
ter cellular retention (Pierré et al. 1990).

Monoclonal antibody (MAb)-antitumour drug
conjugates have gained considerable attention as
potentially useful tools for the treatment of various
human cancers. It is predicted that MABs, selected
specifically to bind particular tumour-associated
surface antigens, will preferentially concentrate the
attached cytotoxic agents at or within a tumour
mass. A number of MAB-vinca alkaloid conjugates
have been synthesised and found to be active in
both *in vitro* and *in vivo* models. A conjugate com-
posed of the adenocarcinoma-reactive MAB KS1/
4S2, coupled through its lysine amino groups to the
4-hydroxy group of the 4-desacetylvinblastine via
a succinate bridge, containing 4 to 6 desacetylvin-
blastine molecules per molecule of KS1/4S2, ex-
hibited a significant antitumour effect *in vivo*
against human lung and colorectal adenocarci-
noma xenografts in nude mice (Spearman et al.
1987). Pharmacokinetic evaluation (in rats and
monkeys) indicated that the conjugate circulated
mostly in the intact form in the plasma and was
catabolised in the liver, with the subsequent ex-
cretion of vinca metabolites into the bile (Marshall
et al. 1987). Recently, other conjugates composed

of a variety of murine MABs coupled to desace-
tylvinblastine hydrazide have been synthesised and
found to exhibit potent antitumour activity *in vivo*
against a number of human solid tumour xeno-
grafts in nude mice, with increased efficacy and
safety compared with unconjugated desacetylvin-
blastine hydrazide or MAB (Lazuzza et al. 1989).

10. Conclusions

The vinca alkaloids continue to be frequently
used as single agents or in combination with other
drugs in routine cancer chemotherapy. Although
their antitumour mechanisms, clinical activity and
toxicity have been well defined, mechanisms of drug
resistance, nature of transport, and metabolic path-
ways require further investigation. Some studies on
MDR in either a clinical setting (Cantwell et al.
1988) or a clinically relevant model (Louie et al.
1986) have led to the conclusion that classic MDR
was not a major mechanism of clinical drug re-
sistance. It may therefore be desirable to extend
cytotoxicity data surveys rather than to remain
limited to gene expression. As drug resistance and
metabolism are cell detoxification processes, it
would be of interest to study the relationship be-
tween the overexpression of the P-glycoprotein and
the possible amplification of cytochrome P-450
isozymes responsible for the metabolism of
vinca alkaloids.

References

- Beck-Hansen NT, Till JE, Ling V. Pleiotropic phenotype of col-
chicine-resistant CHO cells: cross-resistance and collateral sen-
sitivity. *Journal of Cell Physiology* 88: 23-32, 1976
- Beck WT. Vinca-alkaloid resistant phenotype in cultured human
leukemic lymphoblasts. *Cancer Treatment Reports* 67: 875-882,
1983
- Beck WT, Cirtain MC. Continued expression of vinca alkaloid
resistance by CCRF-CEM cells after treatment with tunica-
mycin or pronase. *Cancer Research* 42: 184-189, 1992
- Beck WT, Mueller TJ, Tanzer LR. Altered surface membrane
glycoproteins in vinca alkaloid-resistant human leukemic
lymphoblasts. *Cancer Research* 39: 2070-2076, 1979
- Beck WT, Danks MK, Cirtain MC, van Heiningen JN. Cross-
resistance patterns and antigen expression in vinca alkaloid-
and other multi drug-resistant human leukemic cell lines.
Progress in Clinical and Biological Research 233: 2333-2340,
1986
- Biedler JL, Peterson RHF. Altered plasma membrane glycocon-
jugates of Chinese hamster cells with acquired resistance to

- actinomycin D, daunorubicin and vincristine. In Sartorelli et al. (Eds) *Molecular actions and targets for cancer chemotherapeutic agents*, pp. 453-482. Academy Press, New York, 1981
- Binet S, Chincieu E, Feliou A, Lataste H, Krikorian A, et al. Immunofluorescence study of the action of navelbine, vincristine and vinblastine on mitotic and axonal microtubules. *International Journal of Cancer* 46: 262-266, 1990
- Bleyer WA, Frisby SA, Oliverio VT. Uptake and binding of vincristine by murine leukemia cells. *Biochemical Pharmacology* 24: 633-639, 1975
- Bloemhof H, Van Jijk KN, De Graaf SSN, Vondrig DEMM, Uges DRA. Sensitive method for the determination of vincristine in human serum by high-performance liquid chromatography after on-line column-extraction. *Journal of Chromatography - Biomedical Applications* 572: 171-179, 1991
- Bodey GP, Yap HY, Yap BS, Valdivieso M. Continuous infusion of vindesine in solid tumors. *Cancer Treatment Reviews* 7: 39-47, 1980
- Boré P, Rahmani R, van Cantfort J, Focan C, Cano JP. Pharmacokinetics of a new anticancer drug, navelbine, in patients: comparative study of radioimmunologic and radioactive determination methods. *Cancer Chemotherapy and Pharmacology* 23: 247-251, 1989
- Bosmann HB. Mechanism of cellular drug resistance. *Nature* 233: 566-569, 1971
- Bowman LC, Houghton JA, Houghton PJ. GTP influences the binding of vincristine in human tumor cytosols. *Biochemical and Biophysical Research Communications* 135: 695-700, 1986
- Brade WP. Critical review of pharmacology, toxicology, pharmacokinetics of vincristine, vindesine, vinblastine. *Beiträge zur Onkologie 6. Proceedings of the International Vincaalkaloid Symposium - Vindesine*: 95-123, 1981
- Budman DR, Shulman P, Marks M, Vinciguerra V, Weisberg L, et al. Phase I trial of vinzolidine. *Cancer Treatment Reports* 68: 979-982, 1984
- Cano JP, Rahmani R, Fabre G, Richard B, Lacarelle B, et al. Human hepatocytes as an alternative model to the use of animal in experiments. In Guillouzo (Ed.) *Liver cells drugs*, Vol. 154, pp. 301-307, Colloque INSERM/John Libbey Eurotext Ltd, 1988
- Cantwell BM, Bozzino JM, Corris P, Harris AL. The multidrug resistant phenotype in clinical practice: evaluation of cross resistance to ifosfamide and mesna after VP 16-213, doxorubicin and vincristine (VPAV) for small cell lung cancer. *European Journal of Cancer and Clinical Oncology* 24: 123-129, 1988
- Carlsen SA, Till JE, Ling V. Modulation of membrane drug permeability in Chinese hamster ovary cells. *Biochimica Biophysica Acta* 455: 900-912, 1976
- Cersosimo RJ, Bromer R, Licciardello JT, Kij H. Pharmacology, clinical efficacy and adverse effects of vindesine sulfate, a new vincaalkaloid. *Pharmacotherapy* 3: 259-274, 1983
- Cros S, Wright M, Morimoto M, Lataste H, Couzinier JP, et al. Experimental antitumor activity of navelbine. *Seminars in Oncology* 16 (Suppl. 4): 15-20, 1989
- Dalton WS, Durie BGM, Alberts DS, Gerlach JH, Gress AG. Characterization of a new drug-resistant human myeloma cell line that expresses P-glycoprotein. *Cancer Research* 46: 5125-5130, 1986
- Danks MK, Metzger DW, Ashmun RA, Beck WT. Monoclonal antibodies to glycoproteins of vinca alkaloid-resistant human leukemic cells. *Cancer Research* 45: 3220-3224, 1985
- Dano K. Active outward transport of daunomycin in resistant Ehrlich ascites tumor cells. *Biochimica Biophysica Acta* 323: 466-483, 1973
- De Bruijn MHL, Van Der Blik AM, Biedler JL, Borst P. Differential amplification and disproportionate expression of five genes in three multidrug-resistant Chinese hamster lung cell lines. *Molecular and Cellular Biology* 6: 4717-4722, 1986
- De Smet M, Van Belle SJP, Storme GA, Massart DL. High performance liquid chromatographic determination of vinca alkaloids in plasma and urine. *Journal of Chromatography - Biomedical Applications* 345: 309-321, 1985
- Ferguson PJ, Cass CE. Differential cellular retention of vincristine and vinblastine by cultured human promyelocytic leukemia HL-60/C1 cells/the basis of differential toxicity. *Cancer Research* 45: 5480-5488, 1985
- Ferguson PJ, Phillips JA, Selner M, Cass CE. Differential activity of vincristine and vinblastine against cultured cells. *Cancer Research* 44: 3307-3312, 1984
- Focan C, Mazy V, Lévi F, Bruguerolle B, Zhou XJ, et al. Circadian-varying plasma concentration of vindesine during 48-hour continuous infusion at a constant rate. In Ensminger et al. (Eds) *Update in drug delivery systems*, pp. 291-285, Futura Publishing Co., Inc., Mount Kisco, NY, 1989
- Ford JM, Hait WN. Pharmacology of drugs that alter multidrug resistance in cancer. *Pharmacological Reviews* 42: 155-199, 1990
- Fry DC, Kuby SA, Mildvan AS. ATP-binding site of adenylate kinase: mechanistic implications of its homology with ras-encoded p21, F1-ATPase, and other nucleotide-binding proteins. *Proceedings of the National Academy of Sciences of United States of America* 83: 907-911, 1986
- Haber H, Norris MD, Kavallaris M, Bell DR, Davey RA, et al. Atypical multidrug resistance in a therapy-induced drug-resistant human leukaemic cell line (LALW-2) resistance to vinca alkaloids independent of P-glycoprotein. *Cancer Research* 49: 5281-5287, 1989
- Hacker MP, Dank JR, Ershler WB. Vinblastine pharmacokinetics measured by a sensitive enzyme-linked immunosorbent assay. *Cancer Research* 44: 478-481, 1984
- Hande K, Gay J, Guber J, Greco FA. Toxicity and pharmacology of bolus vindesine injection and prolonged vindesine infusion. *Cancer Treatment Reviews* 7: 25-31, 1980
- Higgins CF, Hiles ID, Salmond GPC, Gill DR, Downie JA, et al. A family of related ATP-binding subunits coupled to many biological processes in bacteria. *Nature* 323: 448-450, 1986
- Houghton PJ, Houghton JA, Bowman LC, Hazelton BE. Therapeutic selectivity of vinca alkaloids: a role for guanosine 5'-triphosphate. *Anti-Cancer Drug Design* 2: 165-179, 1987
- Huhtikangas A, Lehtola T, Lapinjoki S, Lounasmaa M. Specific radioimmunoassay for vincristine. *Planta Medica* 53: 85-87, 1987
- Inaba M, Johnson RK. Decreased retention of actinomycin D as the basis for cross-resistance in anthracycline-resistant sublines of P388 leukemia. *Cancer Research* 37: 4629-4634, 1977
- Jackson DV, Sethi VS, Spurr CL, White DR, Richard FI, et al. Pharmacokinetics of vincristine infusion. *Cancer Treatment Reports* 65: 1043-1048, 1981
- Jehl F, Eebs J, Herlin C, Quiox E, Caillon C, et al. Determination of navelbine and desacetylnavelbine in biological fluids by high-performance liquid chromatography. *Journal of Chromatography - Biomedical Applications* 525: 225-233, 1990
- Jehl F, Quiox E, Leveque D, Pauli C, Breillout F, et al. Pharmacokinetics and preliminary metabolic fate of navelbine in humans as determined by high performance liquid chromatography. *Cancer Research* 51: 2073-2076, 1991
- Johnson IS, Armstrong JG, Gorman M, Burnett JP. The vinca alkaloids: a new class of oncolytic agents. *Cancer Research* 23: 1390-1427, 1963
- Jordan MA, Himes RH, Wilson L. Comparison of the effects of vinblastine, vincristine, vindesine and vinepidine on microtubule dynamics and cell proliferation in vitro. *Cancer Research* 45: 2741-2747, 1985
- Juliano RL, Ling V. A surface glycoprotein modulating drug permeability in Chinese hamster ovary cell mutants. *Biochimica Biophysica Acta* 455: 152-162, 1976
- Kartner N, Riordan JR, Ling V. Cell surface P-glycoprotein associated with multidrug resistance in mammalian cell lines. *Science* 221: 1285-1288, 1983
- Lapinjoki SP, Veräjänkorka H, Lounasmaa M. An enzyme the antineoplastic agent vincristine. *Cancer Research* 48: 113-128, 1986
- Lazussa BC, Nichols CL, Briet al. New antitumor mon- LY203725 and related con- representative in vivo activity 32: 548-555, 1989
- Legha SS. Vincristine neurotoxicity. *Medical Toxicology* 7: 113-128, 1986
- Ling V, Kartner N, Sudo T. S drug-resistance phenotype. *Cancer Treatment Reports* 65: 1043-1048, 1981
- Lothstein L, Hsu SH, Horwi overexpression of two phosphatases with changes in multidrug resistance. *Journal of Biological Chemistry* 264: 10000-10004, 1989
- Louie KG, Hamilton TC, Wu et al. Adriamycin accumulates in cis-sensitive and resistant cells. *Biochemical Pharmacology* 38: 1000-1004, 1989
- Lu K, Yap HY, Loo TL. Clinic by continuous intravenous infusion. *Cancer Research* 49: 5281-5287, 1989
- Mangeney P, Adriamalisoa R. A new class of antitumor derivatives of vinblastine. *Chemistry* 44: 3765-3768, 1984
- Maral R, Bourut C, Chenu E. Cross-resistance of vinca alkaloids and Pharmacology 5: 197-199, 1984
- Maral R, Bourut C, Chenu E. Activity of 5'-nor-anhydrovinorelbine. *Cancer Research* 44: 3765-3768, 1984
- Marshall E, Spearman R, Gooch. Monoclonal antibody-vinca (LY 256787). In Fisher 34. *Metabolism and Disposition* 14: 1000-1004, 1984
- Mathé G, Huijshoven R, Sokal. Phase II clinical trials with biologic malignancies. *Anticancer Research* 12: 115-127, 1992
- Nelson RL. The comparative pharmacokinetics of vindesine, man patients with cancer. *Anticancer Research* 12: 115-127, 1992
- Owells RJ, Donigan DW. Effect of biologic data with phys vinca alkaloids and their efficacy. *Cancer Research* 44: 3765-3768, 1984
- Owells RJ, Hartke CA, Hain. Metabolism of vinblastine in humans. *Cancer Research* 44: 3765-3768, 1984
- Owells RJ, Owens AH, Donigan DW. Vincristine and colchicine. *Physical Research Communications* 12: 1213-1219, 1977a
- Pierré A, Lavielle G, Hauteff. Pharmacological properties of a derivative of vinblastine. *Anticancer Research* 12: 115-127, 1992
- Pontarotti PA, Rahmani R, M. Monoclonal antibodies to anti- amics and kinetics. *Molecular Pharmacology* 38: 1000-1004, 1989
- Rahmani R, Barbet J, Cano JP. Vincristine and vindesine. *Pharmacology* 38: 1000-1004, 1989
- Rahmani R, Bruno R, Fabre. Clinical pharmacokinetic data. *Pharmacology* 38: 1000-1004, 1989
- Rahmani R, Guérin F, Mart

- ic determination of vinca alkaloids in human plasma by reversed-phase HPLC. *Journal of Chromatography* 29: 321, 1985
- cellular retention of vincristine in human promyelocytic leukemia cells. *Cancer Research* 49: 5555-5560, 1989
- Cass CE. Differential activity against cultured cells. *Cancer Research* 49: 5555-5560, 1989
- Croille B, Zhou XJ, et al. Circulation of vindesine during 48-hour constant rate. In Ensminger et al. *Systems*, pp. 291-285, Futura, New York, 1989
- of drugs that alter multidrug resistance. *Clinical Reviews* 42: 155-199, 1990
- ATP-binding site of adenylylation of its homology with ras-nucleotide-binding proteins. *Academy of Sciences of United States of America* 86: 1000-1004, 1989
- M, Bell DR, Davey RA, et al. In a therapy-induced drug-resistance (LALW-2) resistance to vinca alkaloids. *Cancer Research* 49: 5555-5560, 1989
- Vinblastine pharmacokinetics-linked immunosorbent assay. *Cancer Research* 49: 5555-5560, 1989
- A. Toxicity and pharmacology prolonged vindesine infusion. *Cancer Research* 49: 5555-5560, 1989
- C, Gill DR, Downie JA, et al. ng subunits coupled to many. *Nature* 323: 448-450, 1986
- man LC, Hazelton BJ. Therapeutic: a role for guanosine 5'-phosphate. *Design* 2: 165-179, 1987
- oki S, Lounasmaa M. Specific binding. *Planta Medica* 53: 85-87, 1987
- retention of actinomycin D as anthracycline-resistant sublines. *Arch* 37: 4629-4634, 1977
- White DR, Richard FI, et al. e infusion. *Cancer Treatment Reports* 67: 869-874, 1983
- Caillon C, et al. Determination of vindesine in biological fluids by high-performance liquid chromatography. *Journal of Chromatography* 525: 225-233, 1990
- ili C, Breillout F, et al. Pharmacokinetic fate of navelbine in tumor-bearing mice. *Pharmacokinetics and Therapeutics* 41: 61-67, 1987
- an M, Burnett JP. The vinca alkaloids. *Cancer Research* 23: 5555-5560, 1989
- Comparison of the effects of vindesine and vinorelbine on microtubule polymerization in vitro. *Cancer Research* 49: 5555-5560, 1989
- protein modulating drug per-vary cell mutants. *Biochimica et Biophysica Acta* 1976
- Cell surface P-glycoprotein expression in mammalian cell lines. *Cancer Research* 49: 5555-5560, 1989
- Lapinjoki SP, Veräjänkorpä HM, Huhtikangas AE, Lehtola TJ, Lounasmaa M. An enzyme-linked immunosorbent assay for the antineoplastic agent vincristine. *Journal of Immunology* 133: 128, 1986
- Lazurza BC, Nichols CL, Briggs SL, Cullinan GJ, Johnson DA, et al. New antitumor monoclonal antibody-vinca conjugates LY203725 and related compounds: design, preparation, and representative in vivo activity. *Journal of Medicinal Chemistry* 32: 548-555, 1989
- Legha SS. Vincristine neurotoxicity: pathophysiology and management. *Medical Toxicology* 1: 421-427, 1986
- Ling V, Kartner N, Sudo T, Siminovich L, Riordan JR. Multidrug-resistance phenotype in Chinese hamster ovary cells. *Cancer Treatment Reports* 67: 869-874, 1983
- Lothstein L, Hsu SH, Horwitz SB, Greenberger LM. Alternate overexpression of two phosphoglycoprotein genes is associated with changes in multidrug resistance in a J774.2 cell line. *Journal of Biological Chemistry* 264: 16054-16058, 1989
- Louie KG, Hamilton TC, Winker MA, Behrens BC, Tsuruo T, et al. Adriamycin accumulation and metabolism in adriamycin-sensitive and resistant human ovarian cancer cells lines. *Biochemical Pharmacology* 35: 467-472, 1986
- Lu K, Yap HY, Loo TL. Clinical pharmacokinetics of vinblastine by continuous intravenous infusion. *Cancer Research* 43: 1405-1408, 1983
- Mangeney P, Adriamialiso RZ, Langlois N, Langlois Y, Potier P. A new class of antitumor compounds: 5' nor and 5', 6 seco derivatives of vinblastine-type alkaloids. *Journal of Organic Chemistry* 44: 3765-3768, 1979
- Maral R, Bourut C, Chenu E, Mathé G. Experimental in vivo cross-resistance of vinca alkaloid drugs. *Cancer Chemotherapy and Pharmacology* 5: 197-199, 1981
- Maral R, Bourut C, Chenu E, Mathé G. Experimental antitumor activity of 5'-nor-anhydrovinblastine, navelbine. *Cancer Letters* 22: 49-54, 1984
- Marshall E, Spearman R, Goodwin M, Kau D. Disposition of the monoclonal antibody-vinca alkaloid conjugate, KS1/2-DAVLB (LY 256787), in Fisher 344 rats and rhesus monkeys. *Drug Metabolism and Disposition* 15: 640-647, 1987
- Mathé G, Huithoven R, Sokai G, Bayssas M, Beipomme D, et al. Phase II clinical trials with vindesine in patients with hematologic malignancies. *Anticancer Research* 1: 1-9, 1981
- Nelson RL. The comparative clinical pharmacology and pharmacokinetics of vindesine, vincristine and vinblastine in human patients with cancer. *Medical and Pediatric Oncology* 10: 115-127, 1982
- Owells RJ, Donigan DW, Hartke CA, Hains FO. Correlation of biologic data with physicochemical properties among the vinca alkaloids and their congeners. *Biochemical Pharmacology* 26: 1213-1219, 1977a
- Owells RJ, Hartke CA, Hains FO. Pharmacokinetics and metabolism of vinblastine in humans. *Cancer Research* 37: 2597-2602, 1977b
- Owells RJ, Owens AH, Donigan DW. The binding of vinblastine, vincristine and colchicine to tubulin. *Biochemical Biophysical Research Communications* 47: 685-691, 1972
- Pierre A, Lavielle G, Hauteffaye P, Seurre G, Leonce S, et al. Pharmacological properties of a new α -aminophosphonic acid derivative of vinblastine. *Anticancer Research* 10: 139-144, 1990
- Pontarotti PA, Rahmani R, Martin M, Barbet M, Barbet J. Monoclonal antibodies to antitumor vinca alkaloids: thermodynamics and kinetics. *Molecular Immunology* 22: 227-285, 1985
- Rahmani R, Barbet J, Cano JP. A 125 I-radiolabelled probe for vinblastine and vindesine radioimmunoassays. *Clinica Chimica Acta* 129: 57-69, 1983
- Rahmani R, Bruno R, Fabre G, Cano JP. Extrapolation of pre-clinical pharmacokinetic data to therapeutic drug use. *Xenobiotica* 18 (Suppl. 1): 71-88, 1988
- Rahmani R, Guérin F, Martin M, Just S, Cano JP, et al. Comparative pharmacokinetics of antitumor vinca alkaloids: intravenous bolus injections of navelbine and related alkaloids to cancer patients and rats. *Cancer Chemotherapy and Pharmacology* 16: 223-228, 1986
- Rahmani R, Kleisbauer JP, Cano JP, Martin M, Barbet J. Clinical pharmacokinetics of vindesine infusion. *Cancer Treatment Reports* 69: 839-844, 1985
- Rahmani R, Martin M, Barbet J, Cano JP. Radioimmunoassay and preliminary pharmacokinetic studies in rats of 5'-nor-anhydrovinblastine (navelbine). *Cancer Research* 44: 5605-5613, 1984a
- Rahmani R, Martin M, Favre R, Cano JP, Barbet J. Clinical pharmacokinetics of vindesine: repeated treatments by intravenous bolus injections. *European Journal of Cancer and Clinical Oncology* 29: 1400-1417, 1984b
- Rahmani R, Bruno R, Iliadis A, Favre R, Just S, et al. Clinical pharmacokinetics of the antitumor drug navelbine (5'-nor-anhydrovinblastine). *Cancer Research* 47: 5796-5799, 1987
- Rahmani R, Zhou XJ, Boré P, Van Cantfort J, Focan C, et al. Oral administration of [3 H]navelbine in patients: comparative pharmacokinetics using radioactive and radioimmunologic determination methods. *Anti-Cancer Drugs* 2: 405-410, 1991
- Rahmani R, Zhou XJ, Placidi M, Cano JP. In vivo and in vitro pharmacokinetics and metabolism of vinca alkaloids in rat. I. Vindesine (4-desacetyl-vinblastine 3-carboxamide). *European Journal of Drug Metabolism and Pharmacokinetics* 15: 49-55, 1990
- Rahmani-Jourdheuil D, Placidi M, Carcassonne Y, Rahmani R. Uptake of navelbine, a novel antitumor drug, by rat liver plasma membrane vesicles: evidence of passive diffusion and active transport system. *Journal of Cellular Pharmacology* 2: 9-17, 1991
- Rao KSPB, Collard MP, Trouet A. Vinca-23-oyl amino acid derivatives: as new anticancer agents. *Anticancer Research* 5: 379-386, 1985
- Ratain MJ, Vogelzang NJ. Phase I and pharmacological study of vinblastine by prolonged continuous infusion. *Cancer Research* 46: 4827-4830, 1986
- Ratain MJ, Vogelzang NJ, Sinkula JA. Interpatient and intrapatient variability in vinblastine pharmacokinetics. *Clinical Pharmacology and Therapeutics* 41: 61-67, 1987
- Reiss S, Newton KA, Westbury G. Vinca alkaloids and platelets. *New England Journal of Medicine* 299: 310, 1978
- Riordan JR, Deuchars K, Kartner N, Alon N, Trent J, et al. Amplification of P-glycoprotein genes in multidrug-resistant mammalian cell lines. *Nature* 316: 817-819, 1985
- Sethi VS, Burton SS, Jackson D. A sensitive radioimmunoassay for vincristine and vinblastine. *Cancer Chemotherapy and Pharmacology* 4: 183-187, 1980
- Singer WD, Himes RH. Cellular uptake and tubulin binding properties of four vinca alkaloids. *Biochemical Pharmacology* 43: 545-551, 1992
- Spearman ME, Goodwin RM, Apclgren LD, Bumol TF. Disposition of the monoclonal antibody-vinca alkaloid conjugate KS1/4-DAVLB (LY256787) and free 4-desacetylvinblastine (DAVLB) in tumor-bearing nude mice. *Journal of Pharmacology and Experimental Therapeutics* 241: 695-703, 1987
- Todd GC, Gibson WR, Morton DM. Toxicology of vindesine (deacetyl vinblastine amide) in mice, rats and dogs. *Journal of Toxicology and Environmental Health* 1: 843-850, 1976
- Tsuruo T, Lida H, Tsukagoshi S, Sakurai Y. Overcoming of vincristine resistance in P388 leukemia in vivo and in vitro through enhanced cytotoxicity of vincristine and vinblastine by verapamil. *Cancer Research* 41: 1967-1972, 1981
- Tsuruo T, Lida H, Tsukagoshi S, Sakurai Y. Potentiation of vincristine and adriamycin effects in human hemopoietic tumor cell lines by calcium antagonists and calmodulin inhibitors. *Cancer Research* 43: 2267-2272, 1983
- Van den Berg HW, Desai ZR, Wilson R, Kennedy G, Bridges

- JM, et al. The pharmacokinetics of vincristine in man: reduced drug clearance associated with raised serum alkaline phosphatase and dose-limited elimination. *Cancer Chemotherapy and Pharmacology* 8: 215-219, 1982
- Van der Blic AM, Van der Velde-Koeris T, Ling V, Borst P. Overexpression and amplification of five genes in a multidrug-resistant Chinese hamster ovary cell line. *Molecular and Cellular Biology* 6: 1671-1678, 1986
- Vendrig DEMM, Hothuis JJM, Erdélyi-Toth V, Hulshoff A. Solid phase extraction of vinblastine and vincristine from plasma and urine: variable drug recoveries due to non-reproducible column packings. *Journal of Chromatography - Biomedical Applications* 414: 91-100, 1987
- Vendrig DEMM, Teeuwssen J, Hothuis JJM. Analysis of vinca alkaloids in plasma and urine using high performance liquid chromatography with electro-chemical detection. *Journal of Chromatography - Biomedical Applications* 424: 83-94, 1988a
- Vendrig DEMM, Teeuwssen J, Hothuis JJM. Determination of vinca alkaloids in plasma and urine using ion exchange chromatography on silica gel and fluorescence detection. *Journal of Chromatography - Biomedical Applications* 434: 145-155, 1988b
- Walker JE, Saraste M, Runswick MJ, Gay NJ. Distantly related sequences in the α - and β -subunits of ATP synthase, myosin, kinases and other ATP-requiring enzymes and a common nucleotide binding fold. *EMBO Journal* 1: 945-951, 1982
- Welsh MJ, Dedman JR, Brinkley BR. Tubulin and calmodulin. *Journal of Cellular Biology* 81: 624-634, 1979
- Zhou XJ, Boré F, Monjane S, Zouhir S, Favre R, et al. Pharmacokinetics of navelbine after oral administration in cancer patients. *Cancer Chemotherapy and Pharmacology* 29: 66-70, 1991
- Zhou XJ, Martin M, Placidi M, Cano JP, Rahmani R. In vivo and in vitro pharmacokinetics and metabolism of vinca alkaloids. II. Vinblastine and vincristine. *European Journal of Drug Metabolism and Pharmacokinetics* 15: 323-332, 1990
- Zhou XJ, Zhou-Pan XR, Gauthier T, Lacarelle B, Placidi M, et al. Biotransformation of vindesine by human liver microsomal fractions. *Journal de Pharmacie Clinique (Pharmacocinétique: de la recherche à la clinique)*: 231-235, 1992

Correspondence and reprints: Dr R. Rahmani, INSERM U-278, Faculté de Pharmacie, 27 bd, Jean Moulin, 13385 Marseille, Cédex 5, France.

Drugs 44 (Suppl. 4): 17
0012-6667/92/0400-001
© Adis International L
DRSUP3418

Systemic

H.T. Mouridsen
Department of Onc

Summary

ag
of
an

for
Al
th

ha
do
tai
no

tri
ch
fe
li

re
tri
th
at

Endocrine therapy was introduced nearly and since then severalities have been d type of endocrine th mately one-third of i Chemotherapy wa rapidly developed fr to combination regi chemotherapy were r

Round-Table Discussion

Question: Dr Zhou, you comment that the intensity of cellular accumulation of vinca alkaloids may be related to their liposolubility. Does the liposolubility also influence the rate of efflux of vinca alkaloid from the cell?

Dr X.-J. Zhou: Liposolubility appears to influence both the rate and intensity of influx and efflux of vinca alkaloids.

Question: Does there appear to be any correlation between the member of the *mdr1* gene family (i.e. *mdr1a* or *mdr1b*) amplified in a resistant cell line and the overall pattern of multiple drug resistance?

Dr Zhou: In humans, two P-glycoprotein genes are present (*mdr1* and *mdr3*), whereas there are 3 genes in mice (*mdr1a*, *mdr1b* and *mdr2*). cDNA transfection studies demonstrated that mouse MDR1a, MDR1b and human MDR1 can confer multidrug resistance. In contrast, human MDR3 or mouse MDR2 have, so far, failed to confer multidrug resistance.

Question: How do the pharmacokinetics and antitumour activity of oral vinorelbine compare with those of continuous infusion of vinorelbine?

Dr Zhou: Orally administered vinorelbine appears to retain most of the pharmacokinetic properties, side effects and presumably antitumour activity observed after bolus injection of the drug. Regarding continuous infusion, there are no pharmacokinetic data as yet, and only a few results are available concerning the antitumour activity of vinorelbine when given by this route. A recent study (Izzo et al. 1992) reported that, when administered by continuous infusion, vinorelbine was active against breast cancer. Positive clinical responses were obtained and included 5 complete remissions,

15 partial remissions, 7 minor remissions, 18 disease stabilisations and 22 disease progressions. However, more data are necessary to better characterise the antitumour activity, toxicity and pharmacokinetics of vinorelbine when administered by continuous infusion.

Question: Vinorelbine is a cell-cycle-dependent antimitotic agent. Do you think that short infusions of drug would achieve the clinical goal of maximising tumour cell toxicity and minimising adverse effects better than prolonged continuous infusions?

Dr Zhou: Short infusions of vinorelbine can avoid the very high initial concentrations achieved after bolus injections. In this context, it may be expected that short infusions will lead to fewer adverse effects but similar antitumour activity as compared with bolus injections. Appropriate long-term infusion can maintain a constant effective drug level and may presumably result in optimised antitumour activity.

Question: Dr Mouridsen, what considerations should be given to tumour receptor status when deciding which therapy to use in patients with breast cancer?

Dr H.T. Mouridsen: The response to endocrine therapy in advanced breast cancer is related to the receptor (estrogen or progesterone) status of the tumour. Thus, in receptor-positive tumours the chance of a response to endocrine therapy is approximately 50 to 60%, compared with less than 10% in receptor-negative tumours. Therefore, the estrogen receptor status should be used to select patients for first-line systemic therapy, endocrine therapy or chemotherapy. Patients known to be receptor-positive should be offered endocrine therapy.

Those responding to line endocrine therapy the treatment of cl tumours and for the responsive to first-l

In most cases, it originates from an however, the availability be a 10 to 20% rate positive status to reverse. As a consequence of therapy acceptance based upon anal static tissue.

Question: Are there differences in tumor sites when treating bone, liver, brain)?

Dr Mouridsen: Various sites may vary clear why the efficacy at different sites, but biology, local factors involved. Thus, the response in relation to shown higher responses compared with It has been supposed may inhibit the diffusion cerebrospinal fluid. demonstrated that to play activity in brain the activity observed central nervous system

There are no data systemic therapies in disease. However, metastases in vital organs (CNS), chemotherapy is usually recommended of choice because the possibility for the urgent

Question: What is the conjunction with therapy in breast cancer?

Dr Mouridsen: Re

Those responding should also be offered a second-line endocrine therapy, whereas chemotherapy is the treatment of choice for the receptor-negative tumours and for the receptor-positive tumours unresponsive to first-line endocrine therapy.

In most cases, the definition of receptor status originates from analysis of the primary tumour; however, the available data indicate that there may be a 10 to 20% rate of conversion from receptor-positive status to receptor-negative status and vice versa. As a consequence, whenever feasible, selection of therapy according to receptor status should be based upon analysis of the status in the metastatic tissue.

Question: Are there any additional considerations when treating metastases in specific sites (e.g. bone, liver, brain)?

Dr Mouridsen: The response of metastases in various sites may vary between patients. It is unclear why the efficacy of systemic therapy varies in different sites, but it is plausible that tumour biology, local factors and methodological factors are involved. Thus, the available data on rates of response in relation to metastatic site have generally shown higher response rates in soft tissue metastases compared with visceral and bone metastases. It has been supposed that the blood-brain barrier may inhibit the diffusion of some agents into the cerebrospinal fluid. However, many studies have demonstrated that systemic therapy may also display activity in brain metastases that is similar to the activity observed in metastases outside the central nervous system (CNS).

There are no data to support the use of specific systemic therapies according to site of metastatic disease. However, for patients with extensive metastases in vital organs (e.g. the liver, lungs, or CNS), chemotherapy rather than endocrine therapy is usually recommended as the systemic treatment of choice because this approach offers the best possibility for the urgent relief of symptoms.

Question: What role does radiotherapy have in conjunction with chemotherapy or endocrine therapy in breast cancer?

Dr Mouridsen: Radiotherapy has a definite role

in the treatment of locally recurrent or metastatic breast cancer.

In locally recurrent breast cancer in patients who have not had radiotherapy as part of the primary treatment, and who have no evidence of metastatic disease, radiotherapy is often administered with curative intent to the chest wall and regional lymph nodes.

For a larger proportion of patients with metastatic disease, there is a role for radiotherapy as a local palliative measure, in combination with systemic therapy. Patients with bone metastases giving rise to pain that cannot be controlled by medical analgesic therapy, patients with medullary compression or compression of peripheral nerves, and patients with CNS metastases would be expected to benefit from palliative radiotherapy.

Question: Prof. Marty, as single-agent treatment of advanced breast cancer, how does the tolerability profile of vinorelbine compare with the tolerability profiles of other, more traditional single agent regimens?

Prof. M. Marty: The trend in phase II studies has been to increase doses until clinically significant side effects are observed, and this has been the case with taxol, anthracyclines and topoisomerase I inhibitors. Thus, it becomes difficult to compare the therapeutic index of vinorelbine with those of the more traditional single-agent therapies. The adverse effect profile of vinorelbine is dominated by rapidly reversible noncumulative neutropenia, which is amenable to prevention with colony stimulating factors.

Question: What was the rationale for selecting fluorouracil or doxorubicin as the second agent in vinorelbine combination regimens for the treatment of advanced breast cancer?

Prof. Marty: Fluorouracil and doxorubicin were selected for inclusion in combination therapy with the goals of providing adequate first-line therapy and studying an 'optimal' combination based upon the most active agents, i.e. anthracyclines and vinorelbine.

Question: What are some of the other chemotherapeutic agents currently being used in combination with vinorelbine in studies designed to eval-

uate the efficacy and safety of combination therapy in advanced breast cancer?

Prof. Marty: Other combinations currently being studied include combinations of vinorelbine with the following: fluorouracil and doxorubicin; ifosfamide; and thiotepa.

Question: Dr Cvitkovic, vinorelbine has achieved a good objective response rate for the treatment of inoperable non-small cell lung cancer (NSCLC). Does vinorelbine have a place as adjunctive therapy before or after the surgical removal of NSCLC?

Dr E. Cvitkovic: Chemotherapy before surgery or radiotherapy in patients with locoregionally advanced nonmetastatic NSCLC has been shown to be effective in stage IIIA and IIIB disease, both in open studies and in controlled randomised studies of active combinations of drugs. The experience to date with vinorelbine remains anecdotal, but it appears possible to assume that, as clinical experience with vinorelbine as a component of combination therapies in NSCLC accumulates, this agent will be used preoperatively in these patients.

Regarding postoperative adjuvant treatment, there is very little, if any, evidence that such treatment is beneficial in patients with NSCLC. It is certainly desirable to study the possibility of using vinorelbine as postoperative adjuvant therapy, but only in controlled clinical trials.

Question: How many cycles of vinorelbine as a continuous infusion are generally required to achieve a response in the treatment of breast cancer? Is this any different from the number of intravenous bolus cycles required?

Dr Cvitkovic: Responses have been achieved after 3 or 4 cycles of vinorelbine infusions. This does not differ from the number of cycles of vinorelbine administered by bolus injection. However, the data on bolus injections derived from phase II studies are rather poor.

Question: In phase II studies administering vinorelbine 30 mg/m²/week intravenously, dose adjustment and/or treatment delay was required in 70% of patients. Does this imply that the recommended dosage of 30 mg/m²/week is too high for most patients and, if so, are there any recommen-

dations you would make for initiating therapy in the clinical setting?

Dr Cvitkovic: Only 15 to 25% of patients are able to receive vinorelbine 30 mg/m²/week without interruption, and it appears that a dose of 24 to 26 mg/m²/week would be compatible with outpatient administration. However, the dose used should be active and compatible with combined chemotherapy. The maximal dose that can be used in this setting is highly dependent on the type and stage of disease, characteristics of the patient population, other agents included in the combination, administration of haematopoietic growth factors, and the intention of treatment (palliative or curative). We have observed that vinorelbine 18 mg/m² is beneficial when administered on days 1 and 8 of each 3-week cycle in combination with cisplatin/fluorouracil/folinic acid, and that the results observed with this dose are similar to those observed by other investigators using higher doses. Recent Japanese experience has yielded similar results with vinorelbine 20 mg/m²/week for the treatment of NSCLC.

To summarise, there is a need for further research into the potential, synergistic activity and dose-response relationship of vinorelbine.

Question: Dr Sørensen, what are the usual survival times in treated vs untreated small cell lung cancer (SCLC) and NSCLC?

Dr J.B. Sørensen: The survival time in SCLC was 1.5 to 3 months before the introduction of chemotherapy (Hansen 1987). Since the introduction of combination chemotherapy, the median survival time has increased 5-fold. The 2 key factors predicting survival are pretreatment performance status and extent of disease, and the median survival time is now about 9 months and 14 months in patients with extensive and limited disease, respectively (Østerlind & Andersen 1986).

The survival times in patients with inoperable NSCLC receiving the best supportive care only have been compared with the rates observed in patients receiving combination chemotherapy in several studies. The median survival time ranged from 10 to 17 weeks for patients receiving the best supportive care in 3 recent studies (Quoix et al. 1991; Rapp

et al. 1988; Woods et al. 1988). However, the quality of life because of the chemotherapy regimens is a modest improvement from 25 to 33 weeks of receiving chemotherapy; expectation reached in 2 of these studies (Quoix et al. 1988). However, the quality of life because of the chemotherapy regimens is a modest improvement from 25 to 33 weeks of receiving chemotherapy; expectation reached in 2 of these studies (Quoix et al. 1988). However, the quality of life because of the chemotherapy regimens is a modest improvement from 25 to 33 weeks of receiving chemotherapy; expectation reached in 2 of these studies (Quoix et al. 1988).

Question: Is the use of 5-HT₃-receptor antagonists likely to improve the quality of life in patients with lung cancer receiving intensive regimens?

Dr Sørensen: Neutropenic fevers and colic have been valuable additional effects and myelotoxicity has not been a major problem with respect to increase in survival associated with the status of chemotherapy. However, these new drug regimens, but further evaluation is needed in SCLC treatments, although at a low mental level.

Question: Can you report any adverse effects of vinorelbine that have ever been reported?

Dr Sørensen: A few

for initiating therapy in

to 25% of patients are
e 30 mg/m²/week with-
appears that a dose of 24
be compatible with out-
however, the dose used
patible with combined
al dose that can be used
endent on the type and
istics of the patient popu-
led in the combination,
opoietic growth factors,
ment (palliative or cu-
that vinorelbine 18 mg/
ministered on days 1 and
combination with cis-
acid, and that the results
re similar to those ob-
sors using higher doses.
e has yielded similar re-
1 mg/m²/week for the

a need for further re-
synergistic activity and
of vinorelbine.

what are the usual sur-
ntreated small cell lung
?

survival time in SCLC
re the introduction of
37). Since the introduc-
otherapy, the median
15-fold. The 2 key fac-
pretreatment perform-
isease, and the median
months and 14 months
and limited disease, re-
dersen 1986).

patients with inoperable
upportive care only have
es observed in patients
motherapy in several
al time ranged from 10
eiving the best suppor-
Quoix et al. 1991; Rapp

et al. 1988; Woods et al. 1990). In all 3 studies, a modest improvement in survival duration, ranging from 25 to 33 weeks, was observed for patients receiving chemotherapy; the difference in survival expectation reached a statistically significant level in 2 of these studies (Quoix et al. 1991; Rapp et al. 1988). However, the survival advantage for the treated patients is modest and at the expense of quality of life because of the toxicity induced by the chemotherapy regimens.

Question: Is the introduction of new drugs such as 5-HT₃-receptor antagonists and colony-stimulating factors likely to affect the prognosis in patients with lung cancer, by allowing more intensive regimens?

Dr Sørensen: New drugs such as 5-HT₃-receptor antagonists and colony stimulating factors have been valuable additions to available supportive care, since they have reduced gastrointestinal side effects and myelotoxicity. However, until now there has not been a major effect on treatment outcome with respect to increased response rate or survival associated with these agents. Given the current status of chemotherapy in NSCLC, it is unlikely that these new drugs will result in a better prognosis, but further evaluation of colony-stimulating factors in SCLC might contribute to more active treatments, although research is still at the experimental level.

Question: Can you provide more details of the adverse effects of vinorelbine? Has neurotoxicity ever been reported?

Dr Sørensen: A few cases of neurotoxicity have

been reported (Besenval et al. 1991). Among 26 breast cancer patients treated with vinorelbine 30 mg/m² weekly, one patient had WHO grade 3 peripheral neuropathy, which was reversed when treatment was discontinued. One of 78 patients with NSCLC treated with vinorelbine 30 mg/m²/week developed paralytic ileus. This effect was also reversed completely. In addition, 9 cases of muscle weakness (WHO grade 3) were noted in patients treated with vinorelbine for 3 to 6 months.

References

- Besenval M, Leray F, Delgado FM, Merle S, Krikorian A, et al. Navelbine (vinorelbine) tolerance in non-small-cell lung cancer. In Pierre Fabre Oncologie (Eds) Navelbine (vinorelbine). Update and new trends, pp. 185-194. John Libbey Eurotext, Montrouge, 1991
- Hansen HH. Chemotherapy of small-cell carcinoma: a review. Quarterly Journal of Medicine 63: 275-282, 1987
- Izzo J, Cvitkovic E, Toussaint G, Chabot F, et al. High activity and dose intensity (DI) relationship in advanced breast cancer (ABC) with continuous infusion (CIV) of navelbine. Proceedings of the Second Interface of Clinical and Laboratory Responses to Antitumour Drugs: Drugs and Microtubules, p. 56. Marseilles, 2-4 April 1992
- Østerlind K, Andersen PK. Prognostic factors in small-cell lung cancer: multivariate model based on 778 patients treated with chemotherapy with or without irradiation. Cancer Research 46: 4189-4194, 1986
- Quoix E, Dietermann A, Charbonneau J, Boutin C, Meurice JC, et al. La chimiothérapie comportant du cisplatine est-elle utile dans le cancer bronchique non microcellulaire au stade IV? Résultats d'une étude randomisée. Bulletin du Cancer 78: 341-346, 1991
- Rapp E, Pater JL, Willan A, Cormier Y, Murray N, et al. Chemotherapy can prolong survival in patients with advanced non-small-cell lung cancer - report of a Canadian multicenter randomized trial. Journal of Clinical Oncology 6: 633-641, 1988
- Woods RL, Williams CJ, Levi J, Page J, Bell D, et al. A randomized trial of cisplatin and vindesine versus supportive care only in advanced non-small-cell lung cancer. British Journal of Cancer 61: 608-611, 1990